



Skeletal

Procosa®



Unique joint-support supplement with vitamin C and the InCelligence™ Joint-Support Complex

USANA's Procosa supplement helps maintain and protect joint health with vitamin C and a unique InCelligence Joint-Support Complex made with vegetarian glucosamine and Meriva® bioavailable curcumin.*

THE BENEFITS OF PROCOSA

Healthy joints allow the bones to move together comfortably. However, in some individuals, the cartilage breaks down and becomes thin, resulting in more joint friction, discomfort, and stiffness.

In addition to a healthy diet and maintaining moderate physical activity, the unique blend of the InCelligence Joint-Support Complex (formulated with USANA InCelligence Technology™), vitamin C, and other nutrients can help support optimal joint health and function, so you can maintain an active, independent lifestyle.*

THE SCIENCE OF PROCOSA

Between the bones is a narrow space—the joint cavity. Your bones are capped with cartilage, which gets its lubrication and nutrients from synovial fluid, which also acts like a shock absorber. If synovial fluid and cartilage deteriorate, the spacing between the bones narrows, and eventually the bones will rub together.

Research has shown that joint space can be better maintained in those who take glucosamine. Glucosamine also stimulates production of hyaluronic acid, which is responsible for the lubricating and shock-absorbing properties of synovial fluid.*

Glucosamine is a well-absorbed amino sugar that helps regulate normal cartilage formation by encouraging production of collagen and by acting as a building block for proteoglycans. Proteoglycans are essential for healthy cartilage because they bind the water that lubricates and cushions the joint.*

Meriva® is an innovative phytosome that combines curcumin with a phospholipid, so it passes through cell membranes more easily, making it nearly 30 times more bioavailable than traditional curcumin preparations. A potent antioxidant that helps protect joint cells, Meriva has been shown to support joint comfort and function.*

Together, glucosamine and Meriva make up the InCelligence Joint-Support Complex. In addition to the benefits listed above, there are two key cell-signaling processes the complex is designed to promote:

- Exercise can stimulate the immune system and result in temporary inflammation, which is part of the body's natural recovery mechanism for restoring normal structure and function to muscles and tendons. The InCelligence Joint-Support Complex influences the signaling pathways that modulate these responses to help proactively promote a healthy immune and inflammatory response after exercise.*
- Chondrocytes (cartilage cells) maintain healthy function through a natural cell renewal process in which damaged organelles in joint tissue are broken down and removed through a process called autophagy. However, age or other lifestyle factors may slow this process. The InCelligence Joint-Support Complex helps activate autophagy to renew healthy joint-cell function to preserve healthy articular cartilage and support healthy joint aging.*

Procosa also delivers 225 mg of vitamin C because it promotes collagen synthesis. Collagen is one of the main structural proteins that supports the integrity and strength of ligaments and tendons to maintain joint stability.*

THE USANA DIFFERENCE

The InCelligence Joint-Support Complex in Procosa was created using a unique combination of Meriva bioavailable curcumin with high-quality glucosamine hydrochloride. By blending the hydrochloride form of glucosamine with the potassium and magnesium sulfate in the formula, Procosa provides the same benefits of glucosamine sulfate without using shellfish or other animal-based ingredients. Not only is it one of the few vegetarian formulas available on the market, the addition of vitamin C makes Procosa a highly effective, comprehensive approach to joint health.*

*Meriva® is a trademark of Indena S.p.A.

*These statements have not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure, or prevent any disease.



Available for MyHealthPak™



InCelligence

ITEM #131

DIRECTIONS: TAKE THREE (3) TABLETS DAILY, PREFERABLY WITH MEALS.

SUPPLEMENT FACTS		
SERVING SIZE: 3 TABLETS		
AMOUNT PER SERVING		%DV
VITAMIN C (AS CALCIUM ASCORBATE)	225 mg	375%
MANGANESE (AS MANGANESE GLUCONATE)	5 mg	250%
MAGNESIUM (AS MAGNESIUM SULFATE)	43 mg	11%
POTASSIUM (AS POTASSIUM SULFATE)	94 mg	3%
GLUCOSAMINE HCL (VEGETARIAN)	1500 mg	†
MERIVA® BIOAVAILABLE CURCUMIN COMPLEX	247 mg	†

† DAILY VALUE NOT ESTABLISHED.

OTHER INGREDIENTS: MICROCRYSTALLINE CELLULOSE, MODIFIED CELLULOSE, SOY LECITHIN, CROSCARMELOSE SODIUM, ASCORBYL PALMITATE, DEXTRIN, CALCIUM SILICATE, DEXTROSE, SODIUM CITRATE. CONTAINS SOY.

MERIVA® IS A TRADEMARK OF INDENA S.p.A. KEEP OUT OF REACH OF CHILDREN. CONSULT YOUR PHYSICIAN IF YOU ARE PREGNANT, NURSING, TAKING A PRESCRIPTION DRUG, OR HAVE A MEDICAL CONDITION.

LABORATORY TESTED, QUALITY GUARANTEED. MEETS USP SPECIFICATIONS FOR POTENCY, UNIFORMITY, AND DISINTEGRATION WHERE APPLICABLE.

THERE IS A SAFETY SEAL UNDER THE CAP. DO NOT USE IF THE SEAL IS BROKEN OR MISSING. USANA HEALTH SCIENCES, INC., 3838 W. PARKWAY BLVD., SALT LAKE CITY, UTAH 84120, USA.

STORE BELOW 25° C MADE IN USA

References

- Barnes MJ, Kodicek E. Biological hydroxylations and ascorbic acid with special regard to collagen metabolism. 1972. *Vitam Horm* 30:1-43.
- Bisse E, Epting T, Bell A, Lindinger G, Lang H, Wieland H. Reference values for serum silicon in adults. 2005. *Anal Biochem* 337(1):130-5.
- Braham R, Dawson B, Goodman C. The effect of glucosamine supplementation on people experiencing regular knee pain. 2003. *Br J Sports Med* 37:45-9.
- uomo J, Appendino G, Dern AS, Schneider E, McKinnon TP, Brown MJ, Togni S, Dixon BM. Comparative absorption of a standardized curcuminoid mixture and its lecithin formulation. *J Nat Prod*. 2011 Apr 25;74(4):664-9. Epub 2011 Mar 17.
- Matheson AJ, Perry CM. Glucosamine: a review of its use in the management of osteoarthritis. 2003. *Drugs Aging* 20(14):1041-60.
- Nakamura H, Shibakawa A, Tanaka M, Kato T, Nishioka K. Effects of glucosamine hydrochloride on the production of prostaglandin E2, nitric oxide and metalloproteinases by chondrocytes and synovioocytes in osteoarthritis. *Clin Exp Rheumatol*. 2004 May-Jun;22(3):293-9.
- Pavelka K, Gatterova J, Olejarova M, Machacek S, Giacovelli G, Rovati LC. Glucosamine sulfate use and delay of progression of knee osteoarthritis: a 3-year, randomized, placebo-controlled, double-blind study. 2002. *Arch Intern Med* 162(18):2113-23.
- Poolsup N, Suthisitsang C, Channark P, Kittikuluth W. Glucosamine Long-Term Treatment and the Progression of Knee Osteoarthritis: Systematic Review of Randomized Controlled Trials. 2005. *Ann Pharmacotherapy* 39(6):1080-7.
- Reginster JY, Deroisy R, Rovati LC, Lee RL, Lejeune E, Bruyere O, Giacovelli G, Henrotin Y, Dacre JE, Gossett C. Long-Term effects of glucosamine sulphate on osteoarthritis progression: a randomized, placebo-controlled clinical trial. 2001. *Lancet* 357(9252):251-6.
- Richy F, Bruyere O, Ethgen O, Cucherat M, Henrotin Y, Reginster JY. Structural and symptomatic efficacy of glucosamine and chondroitin in knee osteoarthritis: a comprehensive meta-analysis. 2003. *Arch Intern Med* 163(13):1514-22.
- Thie NM, Prasad NG, Major PW. Evaluation of glucosamine sulfate compared to ibuprofen for the treatment of temporomandibular joint osteoarthritis: a randomized double blind controlled 3 month clinical trial. 2001. *J Rheumatology* 28(6):1347-1355.
- Tinker D, Rucker RB. Role of selected nutrients in synthesis, accumulation, and chemical modification of connective tissue proteins. 1985. *Physiol Rev* 65(3):607-57.
- Zheng, O.S. and Chen, A. (2004) Activation of PPAR gamma is required for curcumin to induce apoptosis and to inhibit the expression of extracellular matrix genes in hepatic stellate cells in vitro. *Biochem J* 384(1): 149-157.
- Chen, A. and Xu, J. (2005) Activation of PPAR(gamma) by curcumin inhibits Moser cell growth and mediates suppression of gene expression of cyclin D1 and EGFR. *Am J Physiol Gastrointest Liver Physiol*. 288(3), G447-456.
- Notarbartolo, M., Poma, P., Perri, D., Dusonchet, L., Cervello, M. and D'Alessandro, N. (2005) Antitumor effects of curcumin, alone or in combination with cisplatin or doxorubicin, on human hepatic cancer cells. Analysis of their possible relationship to changes in NF-kB activation levels and in IAP gene expression. *Cancer Lett*. 224(1), 53-65.
- Xu, Y.X., Pindolia, K.R., Janakraman, N., Chapman, R.A. and Gautam, S.C. (1997) Curcumin inhibits IL1 alpha and TNF-alpha induction of AP-1 and NF-kB DNA-binding activity in bone marrow stromal cells. *Hematopathol Mol Hematol*. 11(1), 49-62.
- Carames, B. (2013). Glucosamine activates autophagy in vitro and in vivo. *Arthritis and Rheumatism*. Volume 65 (7); 1843-1852.

